

The genus *Pelophylax* (Amphibia, Ranidae) in Pakistan: museum collections and possible distribution

Daniel Jablonski¹, Addison Wynn², Rafaqat Masroor³, Theodore Papenfuss⁴, Spartak N. Litvinchuk^{5,6}, Glib Mazepa^{7,8}

¹ Department of Zoology, Comenius University in Bratislava, Mlynská dolina, Ilkovičova 6, 842 15, Bratislava, Slovakia

² Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

³ Zoological Sciences Division, Pakistan Museum of Natural History, Garden Avenue, Shakarparian, Islamabad-44000, Pakistan

⁴ Museum of Vertebrate Zoology, University of California, Berkeley, California, USA

⁵ Institute of Cytology, Russian Academy of Sciences, Tikhoretsky pr. 4, Saint Petersburg, 194064, Russia

⁶ Department of Zoology and Physiology, Dagestan State University, Gadzhiev ul. 43-a, Makhachkala, Dagestan, 336700, Russia

⁷ Department of Ecology and Evolution, Biophore Building, University of Lausanne, 1015, Lausanne, Switzerland

⁸ Department of Ecology and Genetics, Evolutionary Biology, Norbyvägen 18D, 75236, Uppsala, Sweden

<http://zoobank.org/4BFD0532-A6BE-440A-9959-E6573B60FB59>

Corresponding author: Daniel Jablonski (daniel.jablonski@uniba.sk)

Academic editor: Günter Gollmann ♦ Received 24 February 2021 ♦ Accepted 7 June 2021 ♦ Published 27 July 2021

Abstract

We provide the first comprehensive data on the questionable distribution of the genus *Pelophylax* and the family Ranidae from Pakistan. Based on a literature review and two specimens of the genus from Tasp, Panjgur District in Pakistani Balochistan (USNM 26194–95), stored in the Smithsonian National Museum of Natural History, Washington, DC, USA, we discuss the possible occurrence and affiliation of these frogs in the context of Central Asia. Our comparison shows that the nearest records of *Pelophylax* in relation to the Tasp specimens are reported from more than 280 km (air-line) away in Iran and Afghanistan, which are currently separated by hot and mostly desert environments. We suggest that possible surviving populations of this genus may still be present in Balochistan (Rakhshan River) or Khyber Pakhtunkhwa (Kabul River) Provinces of Pakistan. This would, however, need further field investigations.

Key Words

Afghanistan, biogeography, Central Asia, Iran, museum data, *Bufotes*, *Chrysopaa*, *Nanorana*, *Rana*

Introduction

The genus *Pelophylax* Fitzinger, 1843 currently represents 20–21 species and 2–3 hemiclones distributed from northern Africa to the Far East in Asia. They are morphologically very similar and often mentioned as particular species complexes (groups) with unresolved phylogenetic relationships (Akin et al. 2010). Populations from the Middle East and Central Asia are ranked under the *P. ridibundus* group and are geographically associated with the

following species: *P. ridibundus* (Pallas, 1771), *P. bedriagae* (Camerano, 1882), *P. caralitanus* (Arikan, 1988) and *P. terentievi* (Mezhzhherin, 1992). All these taxa have phenotypic, ecological and etiological similarities, yet the ongoing research shows that they will need comprehensive evolutionary-taxonomic re-evaluation (Mazepa et al., in prep.). The members of this group are paraphyletic, containing cryptic taxa (Plötner 2005). Populations of the *ridibundus* group have a continuous distribution and are most abundant in the temperate and subtropical zones of

the Western Palearctic, with a scattered distribution in the southern dry regions in the Middle East where water bodies are available. Overall, they are distributed from eastern Spain to western China, through Central Europe, the Balkans, the Anatolian Peninsula, the Caspian Region and Central Asia (Akin et al. 2010). In the division between Central (Afghanistan, Iran, Turkmenistan, Uzbekistan) and Southern Asia, where the territory of Pakistan falls, three species are currently recognised: *P. bedriagae*, *P. ridibundus* and *P. terentievi* (Wagner et al. 2016; Pesarakloo et al. 2017; Showler 2018; Frost 2021).

Pakistan is a country in the transition zone between the Palearctic and Oriental faunas, which is based on the fact that representatives of (not only) amphibians from both these two zoogeographic regions may be found in the country (Khan 2006; Jablonski et al. 2020, 2021). Currently, the territory of Pakistan contains four families of amphibians: Bufonidae Gray, 1825; Dicroglossidae Anderson, 1871; Megophryidae Bonaparte, 1850; and Microhylidae Günther, 1858 (Masroor 2012; Sarwar et al. 2016; Frost 2021). Whereas Bufonidae contains both Palearctic (*Bufo* Rafinesque, 1815) and Oriental (*Duttaphrynus* Frost et al., 2006; *Firouzophrynus* Safaei-Mahroo & Ghaffari, 2020) genera, other families are represented only by genera with mostly Oriental affiliation (e.g. *Euphlyctis* Fitzinger, 1843; *Minervarya* Dubois, Ohler, Biju, 2001; *Microhyla* Tschudi, 1838; *Scutiger* Theobald, 1868).

Previous reports of the family Ranidae from Pakistan (Khan 2006) are for species currently recognized in the family Dicroglossidae. However, historical data mention the genus *Pelophylax* (formerly *Rana* Linnaeus, 1758), a true member of the family Ranidae, for Pakistan. The first *Pelophylax* record ("*Rana esculenta*, L.") was provided by Blanford in Hubrecht (1880: 621) from the locality Nál (Naal) of central Balochistan [coordinates given by Blanford as 29°30'N, 66°E, however, the correct position should be ~27.682°N, 66.183°E sensu St. John (1889)]. Another specimen was collected in Baghunna (~27.907°N, 66.298°E), "*so badly preserved that doubts about its identity with this or any other species may be entertained*" (Hubrecht 1880). These specimens were part of a small collection of amphibians and reptiles made by "*Dr. C. Duke in April and May 1877*" and deposited in the Leiden Museum (currently known as Naturalis Biodiversity Center, Leiden, The Netherlands; Hubrecht 1880). However, in the digital catalogue, we were able to find only a specimen RMNH. RENA.4024 of *Trapelus agilis* (Olivier, 1807) from that collection. We, moreover, think that both localities belong to *Chrysopaa sternosignata* (Murray, 1885), aquatic species that was described later than Blanford in Hubrecht published their observations. Authors thus could affiliate them erroneously with "*Rana esculenta*".

Later, "*Rana esculenta*, L., var. *ridibunda*, Pallas" was mentioned by Boulenger (1898: 353) for Pakistan from the collections of the BMNH as two collection items (i.e. each in a separate jar): i) a tadpole originating from the area "Quetta to Nushki" (~29.71°N, 66.40°E; loc. 2), yet we were able to recover no traces of this specimen (first mentioned by Boulenger in Aitchison 1889: 105) and

ii) a male and a female, NHMUK 1955838 (= BMNH 1884.11.20.6–7) (Boulenger 1882, 1891, 1920: 89–90).

The *Pelophylax* larva was collected by James E. T. Aitchison, naturalist of the Afghan Boundary Commission, on 22–29 September 1884 (Alam 2009). The following settlements were indicated along the road between Quetta and Nushki: Spangaoli, Gobraik, Khanak, Panj-paj, Chaman and Kaisar (Aitchison 1889). Notably, along this route, Aitchison collected an adult specimen of "*Bufo viridis*" NHMUK 1880497 (= 1886.9.21.128) that was later defined as a paratype of *Bufo* *zugmayeri* (Eiselt and Schmidtler 1973). Possible concerns on larval affiliation to *Pelophylax* (identification made by Boulenger) may be ruled out by the fact that Aitchison's collection also contained larvae of a green toad "*Bufo viridis*, Laur.": "*Numerous Tadpoles from streams at base of Mt. Do-Shakh*" (Boulenger in Aitchison 1889: 105).

The adult frog vouchers came from the locality "*near Sarawan, Baluchistan*" (~28.67°N, 65.53°E; loc. 3) and were deposited by George E. Mason (Boulenger 1891: 380). The origin of some other amphibian specimens obtained from Mason was doubted by Boulenger (1920: 89), while no such doubt was raised for the Sarawan specimens.

All abovementioned records come from north-eastern Balochistan, near the border with Afghanistan (Fig. 1).

Materials and methods

During extensive research on the biogeography and taxonomy of the genus *Pelophylax*, we came across a forgotten collection comprising two specimens of *Pelophylax* originating from south-western Pakistan (Balochistan Province), deposited in the Smithsonian Institution National Museum of Natural History, Washington, DC, USA (USNM). We, therefore, provide important data and confirmation of (at least) the historical presence of these frogs for Pakistan (western Balochistan), supported by well available museum material. We took 14 morphometric characters (Table 1) and detailed photos of both specimens (Figs 2, 3). We also compiled, updated, or even corrected (see localities of vouchers CAS [California Academy of Sciences] 115915, FMNH [Field Museum of Natural History] 161114–16 in Suppl. material 1: Table S1 and literature therein) the available field and museum records of the genus *Pelophylax* from eastern Iran and Afghanistan to present the possible geographical affinity to the closest populations and obtain a better view on the origin of Pakistani populations (see Suppl. material 1: Table S1). Moreover, we attempted to genotype the mitochondrial *ND3* locus of specimens (USNM 26194–95) from Pakistan, as well as vouchers from one of the closest known localities of the genus from Iran (Zabol, MVZ [Museum of Vertebrate Zoology] 243552–63 and MVZ 243612–13). However, due to the age and preserved medium of specimens from Pakistan, we were able to extract DNA and sequences from only four of the Zabol vouchers, namely MVZ 243553–54 and 243612–13 (GenBank Accession Numbers: [MW590982–MW590985](#)).

Table 1. Morphological data of two specimens of *Pelophylax* from Pakistan, stored in the Smithsonian National Museum of Natural History. Morphometric data are presented in millimetres and bilateral characters are presented with the left/right sides.

Measurements	USNM 26194	USNM 26195
	(♂)	(♀)
Total length	78	80
Head length	23	26
Distance between nostrils	4	4
Distance between front end of eye to tip of snout	10	11
Distance between the eyes	11	11
Distance between eye lids	2	2
Width of eye lids	9/9	8/8
Diameter of ear membrane	6/6	5/5
Length of femur	39/39	32/34
Length of tibia	40/41	42/43
Length of small toe	12/12	11/10
Length of metatarsal tubercle	4/4	4/4
Width of the metatarsal tubercle	2/1	2/2
Length of the second shank	20/20	19/19

Results

Two *Pelophylax* specimens (“*Rana ridibunda*”; Figs 2, 3) were collected from Tasp (26.99°N, 64.06°E; Fig. 1, loc. 1), south-western Balochistan, Pakistan, by George E. Mason (London, England) and deposited at the National Museum of Natural History, Smithsonian Institution,

Washington, DC, USA, as an exchange for invertebrates (Fig. 4). The time of their collection is unknown, but before they were catalogued in July 1899 (Fig. 4). Both specimens were adult and have voucher numbers USNM 26194 (male; Fig. 2) and USNM 26195 (female, Fig. 3). Together with the material from Sarawan (BMNH 1884.11.20.6–7; see Introduction), these two specimens represent the only vouchered specimens of the genus from Pakistan. The specimens from Tasp are also the western- and southernmost known records of the genus for Pakistan. Morphological characters, together with the visible colour pattern, correspond to the *P. ridibundus* group. We compared the locality of Tasp with available locality records from Pakistan, Afghanistan and eastern Iran (overall 57 identified localities representing 147 vouchered specimens; Fig. 1 and Suppl. material 1: Table S1) to evaluate their geographic distances and closest locality. The nearest locality is “*Hodar, Sarbaz River*” (MMTT [Muzeye Melli-e Tarikh-e Tabiei, Tehran] 877; Fig. 1, loc. 57; Safaei-Mahroo and Ghaffari 2020) in Iran, approximately 280 km (air-line) from Tasp in Pakistan. Other Iranian and Afghan localities are more than 500 km from Tasp (Fig. 1). We also compared the distances to previously known historical records of the genus in Pakistan mentioned by Blanford in Humbrecht (1880) and




Figure 1. Locality data of the genus *Pelophylax* in eastern Iran and Afghanistan, based on museum voucher specimens and published field records (orange circles) in the context of data from Pakistan (green circles). The nearest localities to Tasp (USNM 26194–95; locality no. 1) are marked with arrows. The question mark represents possible presence of the genus *Pelophylax* in Pakistan on the Kabul River. The dashed green line represent the border between the Palearctic and Oriental zoogeographic realms. For details and locality numbers, see Suppl. material 1: Table S1. The individual in the figure originated from Kunduz Province, Afghanistan, photographed by Sjoerd van Bemmél, BemmélPhotography.



Figure 2. The voucher specimen of *Pelophylax* sp. USNM 26194, adult male. Scale bars correspond to 1 cm. Photo credit Esther M. Langan, modified by Glib Mazepa.




Figure 3. The voucher specimen of *Pelophylax* sp. USNM 26195, adult female. Scale bars correspond to 1 cm. Photo credit Esther M. Langan, modified by Glib Mazepa.

CURRENT No.	ORIGINAL No.	NAME	SEX	LOCALITY	WHEN COLLECTED	NATURE OF ORIENT.	CURRENT No.	MEASUREMENT	RECEIVED FROM	COLLECTED BY	CONT.	WHEN EXTENDED	No. of Specimens	REMARKS
2618	1	<i>Phrynosoma hernandesi</i>		Michouding Co.			1	35278	Wm. D. Brown			14	1	PARATYPE
	2	<i>Phrynosoma hernandesi</i>		Michouding Co.			2	35286	V. R. Bodley				1	
	3	<i>Phrynosoma hernandesi</i>		Michouding Co.			3	35284	Wm. D. Brown			15	1	
	4	<i>Phrynosoma hernandesi</i>		Michouding Co.			4						2	Egg with young
	5	<i>Rana catesbeiana</i>		Michouding Co.	Aug 5 1894		5	35238	Geo. E. Mason	Geo. E. Mason		18	1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
2619	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
2620	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5	"		"	"		5						1	
	6	"		"	"		6						1	
	7	"		"	"		7						1	
	8	"		"	"		8						1	
	9	"		"	"		9						1	
	0	"		"	"		0						1	
	1	"		"	"		1						1	
	2	"		"	"		2						1	
	3	"		"	"		3						1	
	4	"		"	"		4						1	
	5</													

tunkhwa Province in the north-western part of the country (Fig. 1). The Kabul River is a biogeographic corridor for the Oriental fauna into the Palearctic area that probably allowed colonisation of valleys of Hindu Kush mountains around the river from Pakistan to eastern Afghanistan [e.g. *Euphlyctis cyanophlyctis* (Schneider, 1799); *Calotes versicolor* (Daudin, 1802); *Ophisops jerdonii* Blyth, 1853; *Varanus bengalensis* (Daudin, 1802); *Fowlea piscator* (Schneider, 1799); Wagner et al. 2016; Jablonski and Lesko 2018; Jablonski et al. 2019]. Given the records of *Pelophylax* in Afghanistan, for example, from Paghman near Kabul (FMNH: 161055; Anderson and Leviton 1969; Fig. 1 loc. 5) and especially from the locality “ca. 25 km SE (by air) Jalalabad” (MVZ: 236876; Fig. 1 loc. 4), only 60 km from the Pakistani border, such vice versa colonisation and presence of the genus *Pelophylax* in north-western Pakistan is highly possible. In this context, we checked the specimens stored in the Florida Museum of Natural History at the University of Florida (UF) and vouchered as UF 82322–24, collected in Yakh Tangi, Khyber Pakhtunkhwa Province, Pakistan (approximately 34.30°N, 71.65°E). These specimens were stored in the collection under the name “*Rana* “*farooqi*” nov. sp. Auffenberg”, raising serious questions on the identification of these specimens as *Pelophylax*. However, available photos reveal that these specimens might be members of the genus *Nanorana* Günther, 1896, most likely *N. vicina* (Stoliczka, 1872).

Acknowledgements

We would like to thank E. Ely (Californian Academy of Sciences, San Francisco, California, USA); C. Spencer (Museum of Vertebrate Zoology, Berkeley, California, USA) for assistance with information regarding collection and loan of the *Pelophylax* tissue samples from Zabol, eastern Iran; R. Wilson (Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, USA) for the loan of tissue samples of *Pelophylax* from Pakistan; E. M. Langan (Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, USA) for the photographs of USNM 26194 and 26195; C. M. Sheehy III and D. Blackburn (Florida Museum of Natural History at the University of Florida, USA) for information and photos of “*Rana farooqi*”; and S. van Bommel (The Netherlands) for his photo of *P. terentievi* from Afghanistan. We also thank two reviewers for their useful comments. This work was supported to DJ by the Slovak Research and Development Agency under the contract APVV-19-0076 and by the grant of the Scientific Grant Agency of the Slovak Republic VEGA 1/0286/19.

References

- Aitchison JET (1889) III. The Zoology of the Afghan Delimitation Commission. Transactions of the Linnean Society of London, 2nd Series 5: 53–142. <https://doi.org/10.1111/j.1096-3642.1889.tb00159.x>
- Akin C, Bilgin CC, Beerli P, Westaway R, Ohst T, Litvinchuk SN, Uzzel T, Bilgin M, Hotz H, Guex G-D, Plötner J (2010) Phylogeographic patterns of genetic diversity in eastern Mediterranean water frogs have been determined by geological processes and climate change in the Late Cenozoic. Journal of Biogeography 37(11): 2111–2124. <https://doi.org/10.1111/j.1365-2699.2010.02368.x>
- Alam M (2009) Plant collectors in Afghanistan. Bulletin de la Société vaudoise des Sciences naturelles 91: 301–340.
- Anderson SC, Leviton AE (1969) Amphibians and reptiles collected by the Street expedition to Afghanistan, 1965. Proceedings of the California Academy of Sciences, Fourth Series 37(2): 25–56.
- Blanford WT, Hubrecht AAW (1880) On a collection of reptiles and amphibians from Baluchistan made by Dr. C. Duke in April and May, 1877. With a note by W.T. Blanford, F.Z.S. Proceedings of the Zoological Society of London 1880: 621.
- Boulenger GA (1882) Catalogue of the Batrachia Salientia s. Ecaudata in the collections of the British Museum. London, 256 pp.
- Boulenger GA (1889) Reptiles and batrachians. III. 94–106. In: Aitchison JET (Ed.) The Zoology of the Afghan Delimitation Commission. Transactions of the Linnean Society of London, 2nd Series. <https://doi.org/10.1111/j.1096-3642.1889.tb00162.x>
- Boulenger GA (1891) A contribution to the knowledge of the races of *Rana esculenta* and their geographical distribution. Proceedings of the Zoological Society of London 1891: 374–384. <https://doi.org/10.1111/j.1096-3642.1891.tb01762.x>
- Boulenger GA (1898) The tailless batrachians of Europe. London V2. 376 pp. <https://www.biodiversitylibrary.org/page/36107722>
- Boulenger GA (1920) A monograph of the South Asian, Papuan, Melanesian and Australian frogs of the genus *Rana*. Records of the Indian Museum 20: 1–226. <https://www.biodiversitylibrary.org/page/12748884>
- Frost DR (2021) Amphibian Species of the World: an Online Reference. Version 6.1 (28 Januar 2021). Electronic Database accessible at <https://amphibiansoftheworld.amnh.org/index.php>. American Museum of Natural History, New York. <https://doi.org/10.5531/db.vz.0001>
- Jablonski D, Lesko AJ (2018) New locality record of the Bengal monitor, *Varanus bengalensis* (Daudin, 1802), from Afghanistan. Herpetology Notes 11: 915–917.
- Jablonski D, Khan MA, Masroor R (2020) The genus *Microhyla* (Anura: Microhylidae) in Pakistan: species status and origins. Zootaxa 4845: 293–296. <https://doi.org/10.11646/zootaxa.4845.2.11>
- Jablonski D, Masroor R, Hofmann S (2021) Revisited molecular phylogeny of the genus *Sphaerothera* (Anura: Dicroglossidae): the biogeographic status of northernmost populations and further taxonomic changes. Diversity 13: 216. <https://doi.org/10.3390/d13050216>
- Jablonski D, Regan JM, Holzheuser C, Farooqi J, Basit A, Masroor R (2019) Additional data to the herpetofauna of Afghanistan. Herpetozoa 32: 177–193. <https://doi.org/10.3897/herpetozoa.32.e38171>
- Khan MS (2006) Amphibians and reptiles of Pakistan. Krieger Publishing Company, Malabar, Florida, 311 pp.
- Masroor R (2012) A contribution to the herpetology of northern Pakistan: The amphibians and reptiles of Margalla Hills National Park and surrounding regions. Society for the Study of Amphibians and Reptiles (SSAR), Ithaca, New York, USA and Chimaira Buchhandelsgesellschaft mbH, Germany, 217 pp.
- Nicolas V, Mataame A, Crochet P-A, Geniez P, Ohler A (2015) Phylogeographic patterns in North African water frog *Pelophylax sahariensis* (Anura: Ranidae). Journal of Zoological Systematics and Evolutionary Research 53: 239–248. <https://doi.org/10.1111/jzs.12094>

- Pesarakloo A, Rastegar-Pouyani E, Rastegar-Pouyani N, Kami H, Khosravani A, Oraie H (2017) The first taxonomic reevaluation of the Iranian water frogs of the genus *Pelophylax* (Anura: Ranidae) using sequences of the mitochondrial genome. *Mitochondrial DNA* 28: 392–398. <https://doi.org/10.3109/19401736.2015.1127362>
- Plötner J (2005) Die westpaläarktischen Wasserfrösche – Von Märtyrern der Wissenschaft zur biologischen Sensation. Laurenti Verlag, Bielefeld, 160 Seiten.
- Safaei-Mahroo B, Ghaffari H (2020) The Complete Guide to Amphibians of Iran: Biology, Ecology, and Conservation. University of Kurdistan Press, 331 pp.
- Sarwar MK, Malik MF, Azam MHI, Iqbal W, Ashiq U (2016) Distribution and current status of amphibian fauna of Pakistan: a review. *Electronic Journal of Biology* 12(3): 243–246.
- Showler DA (2018) A checklist of the Amphibians and Reptiles of the Republic of Uzbekistan with a review and summary of species distribution. <https://www.sustainablehoubaramanagement.org/wp-content/uploads/2018/09/Uzbekistan-Amphibian-Reptile-Checklist-14Sept2018-PDF.pdf> [accessed at March 1, 2020]
- St. John OB (1889) On birds of Southern Afghanistan and Kelát. *Ibis* 1: 145–180. <https://doi.org/10.1111/j.1474-919X.1889.tb06382.x>
- Wagner P, Bauer AM, Leviton AE, Wilms TM, Böhme W (2016) A checklist of the amphibians and reptiles of Afghanistan. *Exploring herpetodiversity using biodiversity archives. Proceedings of the Californian Academy of Sciences* 63: 457–565.

Supplementary material 1

Table S1. An overview of localities and voucher specimens from Afghanistan, eastern Iran, and Pakistan used in this study

Authors: Daniel Jablonski, Addison Wynn, Rafaqat Masroor, Theodore Papenfuss, Spartak N. Litvinchuk, Glib Mazepa

Data type: species data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.34.64955.suppl1>